

3 Equatorial Wave Theory

Lecture based on 'An Introduction to Dynamic Meteorology' by J. R. Holton, *Academic Press, INC.*, 3rd edition, 511 pp.




Figure 7: Dispersion diagram for tropical Outgoing longwave radiation. Source: www.cgd.ucar.edu, NOAA.

Equatorial waves are an important class of eastward and westward propagating disturbances that are trapped about the equator (that is, they decay away from the equatorial region). In a dispersion diagram for observed equatorial quantities, these wave can be identified as regions of increased energy density (Fig. 7).

Diabatic heating by organized tropical convection may excite equatorial wave motions (see Fig. 8). Through such waves the dynamical effects of convective storms can be communicated over large longitudinal distances in the tropics. Such waves can produce remote responses to localized heat sources. Furthermore, by influencing the pattern of low-level moisture convergence they can partly control the spatial and temporal distribution of convective heating. In order to introduce equatorial waves in the simplest possible context, we here use a shallow-water model and concentrate

SST-induced heating may induce waves....




Figure 8: Equatorial Kelvin and Rossby waves triggered by an SST-induced heating. Source: Kucharski et al. 2008: A Gill-Matsuno-type mechanism explains the tropical Atlantic influence on African and Indian monsoon rainfall. *Q. J. R. Meteorol. Soc.* (2009), 135, 569-579, DOI: 10.1002/qj.406

on the *horizontal* structure.

3.1 The shallow water equations

The shallow water equations are a drastic simplification to the real atmospheric flow. However, despite its simplicity it gives often a good insight into many atmospheric wave phenomena. The basic assumptions in the shallow water model are

- (i) The flow is incompressible $\rho = \text{const.}$
- (ii) The flow is shallow enough so that the horizontal velocity components are independent of height.
- (iii) The flow is hydrostatic. Accelerations in the vertical direction may be neglected.

Let us consider the horizontal momentum equations 1 and the hydrostatic equation ??

$$\frac{\partial u}{\partial t} + (\mathbf{v} \cdot \nabla)u = -\frac{1}{\rho} \frac{\partial p}{\partial x} + fv \quad (56)$$

$$\frac{\partial v}{\partial t} + (\mathbf{v} \cdot \nabla)v = -\frac{1}{\rho} \frac{\partial p}{\partial y} - fu \quad (57)$$

$$\frac{1}{\rho} \frac{\partial p}{\partial z} = -g, \quad (58)$$

Further consider the continuity equation

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0. \quad (59)$$

Integrating the hydrostatic equation from a height z to the top of the fluid leads to (assuming the pressure is vanishing there)

$$\int_z^{h(x,y,t)} \frac{\partial p}{\partial z} dz = - \int_z^{h(x,y,t)} \rho g dz, \quad \text{or} \quad (60)$$

$$-p(x, y, z, t) = -\rho g [h(x, y, t) - z]. \quad (61)$$

Thus the horizontal pressure gradient force in the equations of motion 56, 57 may be expressed as

$$-\frac{1}{\rho} \frac{\partial p}{\partial x} = -g \frac{\partial h}{\partial x} = -\frac{\partial \Phi}{\partial x} \quad (62)$$

$$-\frac{1}{\rho} \frac{\partial p}{\partial y} = -g \frac{\partial h}{\partial y} = -\frac{\partial \Phi}{\partial y}, \quad (63)$$

where we have defined $\Phi(x, y, t) = gh(x, y, t)$. Thus, keeping in mind that there the horizontal velocities do not depend on the vertical direction and ignoring the coriolis term proportional to the vertical velocity, the horizontal equations of motion may be written as

$$\frac{\partial u}{\partial t} + (\mathbf{v} \cdot \nabla)u = -\frac{\partial \Phi}{\partial x} + fv \quad (64)$$

$$\frac{\partial v}{\partial t} + (\mathbf{v} \cdot \nabla)v = -\frac{\partial \Phi}{\partial y} - fu, \quad (65)$$

The number of dependent variables in Eqs. 64 and 65 is reduced to 3, (u, v, Φ) . Thus, if we have another equation only containing (u, v, Φ) , then the system may be complete. This is achieved by simplification of the continuity equation 59 and vertical integration. First, we note that because of $\rho = \text{const}$, Eq. 59 reduces to

$$\frac{\partial w}{\partial z} = -\nabla \cdot \mathbf{v}. \quad (66)$$

If we integrate this equation vertically from 0 to $h(x, y, t)$ we have

$$\int_0^h \frac{\partial w}{\partial z} dz = - \int_0^h \nabla \cdot \mathbf{v} dz \quad (67)$$

$$w(h) := \frac{dh}{dt} = \frac{\partial h}{\partial t} + \mathbf{v} \cdot \nabla h = -(\nabla \cdot \mathbf{v})h \quad (68)$$

Eq. 68 may as well be written as

$$\frac{\partial \Phi}{\partial t} + \mathbf{v} \cdot \nabla \Phi = -\Phi \nabla \cdot \mathbf{v} . \quad (69)$$

Eqs. 64, 65 and 69 build a complete set of differential equations for (u, v, Φ) , and are called the *shallow water equations*.

3.2 Linearization for an Equatorial β -plane

Now we linearize the set of equations 64, 65 and 69 about a motionless mean state with height h_e on an equatorial β -plane. Generally speaking, the β -plane assumption states that $f = 2|\omega| \sin \phi \approx f_0 + \beta y$, that is the $\sin \phi$ -dependence is approximated linearly for a given latitude ϕ_0 by a Taylor series expansion (therefore $\beta = 2|\omega| \cos \phi_0 / a$; a being the mean radius of the earth). If we set the base point at the equator we have $f_0 = 0$, therefore $f \approx \beta y$.

$$\frac{\partial u'}{\partial t} = -\frac{\partial \Phi'}{\partial x} + \beta y v' \quad (70)$$

$$\frac{\partial v'}{\partial t} = -\frac{\partial \Phi'}{\partial y} - \beta y u' \quad (71)$$

$$\frac{\partial \Phi'}{\partial t} = -g h_e \left(\frac{\partial u'}{\partial x} + \frac{\partial v'}{\partial y} \right) , \quad (72)$$

where the primed variables denote the perturbations from the basic state. This is our basic set of linearized equations (with variable coefficients!) to study equatorial wave dynamics. By adjusting the scale height h_e as well the ocean case may be included.

Discuss Inertia-Gravity waves for extratropical situation and approximation $f = f_0 = \text{const}$, and assume $u'(x, t), v'(x, t), \Phi'(x, t)$.

3.2.1 Equatorial Rossby and Rossby-Gravity Modes

In order to find solutions to the linearized system 70, 71 and 72, we assume that the y -dependence can be separated

$$\begin{pmatrix} u' \\ v' \\ \Phi' \end{pmatrix} = \begin{bmatrix} \hat{u}(y) \\ \hat{v}(y) \\ \hat{\Phi}(y) \end{bmatrix} e^{i(kx - \nu t)} . \quad (73)$$

Substitution of Eq. 73 into 70-72 then yields a set of ordinary differential equations in y for the meridional structure functions \hat{u} , \hat{v} , $\hat{\Phi}$:

$$-i\nu\hat{u} = -ik\hat{\Phi} + \beta y\hat{v} \quad (74)$$

$$-i\nu\hat{v} = -\frac{\partial\hat{\Phi}}{\partial y} - \beta y\hat{u} \quad (75)$$

$$-i\nu\hat{\Phi} = -gh_e \left(ik\hat{u} + \frac{\partial\hat{v}}{\partial y} \right). \quad (76)$$

If Eq. 74 is solved for $\hat{u} = k/\nu\hat{\Phi} + i\beta y\hat{v}/\nu$ and inserted into Eq. 75 and 76 we obtain

$$(\beta^2 y^2 - \nu^2)\hat{v} = ik\beta y\hat{\Phi} + i\nu \frac{\partial\hat{\Phi}}{\partial y} \quad (77)$$

$$(\nu^2 - gh_e k^2)\hat{\Phi} + i\nu gh_e \left(\frac{\partial\hat{v}}{\partial y} - \frac{k}{\nu}\beta y\hat{v} \right) = 0. \quad (78)$$

Finally, Eq. 78 is inserted into Eq. 77 to eliminate $\hat{\Phi}$, yielding a second-order differential equation in the single unknown, \hat{v}

$$\frac{\partial^2\hat{v}}{\partial y^2} + \left[\left(\frac{\nu^2}{gh_e} - k^2 - \frac{k}{\nu}\beta \right) - \frac{\beta^2 y^2}{gh_e} \right] \hat{v} = 0. \quad (79)$$

We seek solutions of this equation for the meridional distribution of \hat{v} , subject to the boundary condition that the disturbance fields vanish for $|y| \rightarrow \infty$. This boundary condition is necessary since the approximation $f \approx \beta y$ is not valid for latitudes much beyond $\pm 30^\circ$, so that solutions must be equatorially trapped if they are to be good approximations to the exact solutions on the sphere. Equation 79 differs from the classic equation for a harmonic oscillator in y because the coefficient in square brackets is not a constant but is a function of y . For sufficiently small y this coefficient is positive and solutions oscillate in y , while for large y , solutions either grow or decay in y . Only the decaying solutions, however, can satisfy the boundary conditions.

It turns out that solutions to Eq. 79 which satisfy the condition of decay far from the equator exist only when the constant part of the coefficient in square brackets satisfies the relationship (which is as well the dispersion relation!)

$$\frac{\sqrt{gh_e}}{\beta} \left(-\frac{k}{\nu}\beta - k^2 + \frac{\nu^2}{gh_e} \right) = 2n + 1; \quad n = 0, 1, 2, \dots \quad (80)$$

which is a cubic dispersion equation determining the frequencies of permitted equatorially trapped free oscillations for zonal wave number k and meridional mode number n . These solutions can be expressed most conveniently if y is replaced by the nondimensional meridional coordinate

$$\xi = \left(\frac{\beta}{\sqrt{gh_e}} \right)^{1/2} y. \quad (81)$$

With the Eqs. 80 and 81, Eq. 79 becomes

$$\frac{\partial^2 \hat{v}}{\partial \xi^2} + (2n + 1 - \xi^2) \hat{v} = 0 . \quad (82)$$

This is the differential equation for a quantum mechanical, simple harmonic oscillator. The solution has the form

$$\hat{v}(\xi) = H_n(\xi) e^{-\xi^2/2} , \quad (83)$$

where $H_n(\xi)$ designates the n th *Hermite polynomial*. The first of these polynomials have the values

$$H_0 = 1, \quad H_1(\xi) = 2\xi, \quad H_2(\xi) = 4\xi^2 - 2 . \quad (84)$$

Thus, the index n corresponds to the number of nodes in the meridional velocity profile in the domain $|y| < \infty$. Inserting the solution 83 into Eq. 82 leads to one of the defining differential equations for Hermite polynomials. In general, the three solutions of Eq. 80 can be interpreted as eastward- and westward-moving equatorially trapped gravity waves and westward-moving equatorial Rossby waves. The case $n = 0$ (for which the meridional velocity perturbation has a gaussian distribution centered at the equator) must be treated separately. In this case the *dispersion relationship* 80 (which is something like a characteristic equation that gives the $\nu(k)$ -dependence from which we may derive the phase velocities) factors as

$$\left(\frac{\nu}{\sqrt{gh_e}} - \frac{\beta}{\nu} - k \right) \left(\frac{\nu}{\sqrt{gh_e}} + k \right) = 0 . \quad (85)$$

The root $\nu/k = -\sqrt{gh_e}$, corresponding to a westward-propagating gravity wave, is not permitted since the second term in parentheses in Eq. 85 was explicitly assumed not to vanish when Eqs. 77 and 78 were combined to eliminate Φ . The roots given by the first term in parentheses in Eq. 85 are

$$\nu = k\sqrt{gh_e} \left[\frac{1}{2} \pm \frac{1}{2} \left(1 + \frac{4\beta}{k^2\sqrt{gh_e}} \right)^{1/2} \right] . \quad (86)$$

The positive root corresponds to an eastward-propagating equatorial inertio-gravity wave, while the negative root corresponds to a westward-propagating wave, which resembles an inertio-gravity wave for long zonal scale $k \rightarrow 0$ and resembles a Rossby wave for zonal scales characteristic of synoptic-scale disturbances. This mode is generally referred to as a Rossby-gravity wave.

3.2.2 Equatorial Kelvin Waves

In addition to the modes discussed in the previous section, there is another equatorial wave that is of great practical importance. For this mode, which is called the equatorial *Kelvin wave*, the meridional velocity perturbation vanishes and Eqs. 74 to 76 are reduced to the simpler set

$$-i\nu\hat{u} = -ik\hat{\Phi} \quad (87)$$

$$\beta y\hat{u} = -\frac{\partial\hat{\Phi}}{\partial y} \quad (88)$$

$$-i\nu\hat{\Phi} = -gh_e(ik\hat{u}) . \quad (89)$$

Eliminating Φ between Eq. 87 and Eq. 89, we see that the Kelvin wave dispersion equation is that of the shallow-water gravity wave

$$c^2 = \left(\frac{\nu}{k}\right)^2 = gh_e . \quad (90)$$

According to Eq. 90 the phase speed c can be either positive or negative. But, if Eq. 87 and Eq. 88 are combined to eliminate Φ we obtain a first-order equation for determining the meridional structure

$$\beta y\hat{u} = -c\frac{\partial\hat{u}}{\partial y} , \quad (91)$$

which may be integrated immediately to yield

$$\hat{u} = u_0 e^{-\beta y^2/(2c)} , \quad (92)$$

where u_0 is the amplitude of the perturbation zonal velocity at the equator. Equation 92 shows that if solutions decaying away from the equator are to exist, the phase speed must be positive ($c > 0$). Thus Kelvin waves are eastward propagating and have zonal velocity and geopotential perturbations that vary in latitude as Gaussian functions centered on the equator. The e-folding decay width is given by

$$Y_K = |2c/\beta|^{1/2} , \quad (93)$$

which for a phase speed $c = 30 \text{ m s}^{-1}$ gives $Y_K = 1600 \text{ km}$. The meridional force balance for the Kelvin mode is an exact geostrophic balance between the zonal velocity and the meridional pressure gradient. It is the change in sign of the Coriolis parameter at the equator that permits this special type of equatorial mode to exist.




Fig.1 Horizontal structure of (a) Kelvin wave and (b) mixed-Rossby gravity wave. Contours show geopotential height component and arrows show horizontal wind components.

Figure 9: Illustration of Kelvin (upper panel) and Rossby-gravity (lower panel) waves.




Figure 10: Dispersion diagram for equatorial Rossby-gravity and Kelvin waves.